

Chapter - 4

Carbon and its Compounds

Introduction :

- The element carbon is non-metal. Its symbol is C.
- Carbon is a versatile element the percentage of carbon present in earth crust in form of mineral is 0.02% and in atmosphere as CO_2 is 0.03%.
- All the living things, plants and animals are made up of carbon based compounds.

Carbon always form covalent bonds :

The atomic number of carbon is 6.


Electronic configuration :

| | | |
|-------|---|---|
| | K | L |
| C (6) | 2 | 4 |

How carbon attain noble gas configuration ?

- Carbon is tetravalent, it does not form ionic bond by either losing four electrons (C^{4+}) or by gaining four electrons (C^{4-}). It is difficult to hold four extra electron and would require large amount of energy to remove four electrons. So, carbon can form bond by sharing of its electrons with the electrons of other carbon atom or with other element and attain noble gas configuration.
- The atoms of other elements like hydrogen, oxygen and nitrogen, chlorine also form bonds by sharing of electrons.
- The bond formed by sharing of electrons between same or different atoms is covalent bond.


(i) H_2



One shared pair of electron

H – H single bond between hydrogen atoms


(ii) O_2



Two shared pair of electron

O = O double bond between oxygen atoms



(iii) N_2



Three shared pair of electrons

$\text{N} \equiv \text{N}$ triple bond between nitrogen atoms

Molecule of water has single covalent bond between one oxygen and two hydrogen atoms.



Physical Properties of Covalent Compounds

- (a) Covalent compounds have low melting and boiling points as they have weak intermolecular force.
- (b) They are generally poor conductor of electricity as electrons are shared between atoms and no charged particles are formed.

Versatile Nature of Carbon

The two characteristic properties of carbon element which lead to the formation of large number of compounds :

- (i) **Catenation** : Carbon can link with carbon atoms by means of covalent bonds to form long chains, branched chains and closed ring compound. Carbon atoms may be linked by single, double or triple bonds.
- (ii) **Tetravalency** : Carbon has 4 valence electrons. Carbon can bond with four carbon atoms, monovalent atoms, oxygen, nitrogen and sulphur.


Saturated and Unsaturated Carbon Compounds

Compounds made up of hydrogen and carbon are called hydrocarbon.

| Hydrocarbon | | |
|---|---|--|
| Saturated | Unsaturated | |
| <ul style="list-style-type: none">Single bond between carbon atoms. | <ul style="list-style-type: none">Double or triple bond between carbon atoms. | |
| <ul style="list-style-type: none">$-C-C-$ | <ul style="list-style-type: none">$-C=C-$ | <ul style="list-style-type: none">$-C\equiv C-$ |
| <ul style="list-style-type: none">Alkanes | <ul style="list-style-type: none">Alkenes | <ul style="list-style-type: none">Alkynes |
| General formulae | | |
| C_nH_{2n+2} | C_nH_{2n} | C_nH_{2n-2} |

Electron Dot Structure of Saturated Hydrocarbons

Ethane C_2H_6




The names, molecular formulae and saturated formulae of saturated hydrocarbons (alkanes) are given below :


| Name of Hydrocarbon | Molecular formula | Structural Formula |
|---------------------|---------------------------|---|
| 1. Methane | CH_4 | $\begin{array}{c} \text{H} \\ \\ \text{H} - \text{C} - \text{H} \\ \\ \text{H} \end{array}$ |
| 2. Ethane | C_2H_6 | $\begin{array}{cc} \text{H} & \text{H} \\ & \\ \text{H} - \text{C} & - \text{C} - \text{H} \\ & \\ \text{H} & \text{H} \end{array}$ |
| 3. Propane | C_3H_8 | $\begin{array}{ccc} \text{H} & \text{H} & \text{H} \\ & & \\ \text{H} - \text{C} & - \text{C} - & \text{C} - \text{H} \\ & & \\ \text{H} & \text{H} & \text{H} \end{array}$ |
| 4. Butane | C_4H_{10} | $\begin{array}{cccc} \text{H} & \text{H} & \text{H} & \text{H} \\ & & & \\ \text{H} - \text{C} & - \text{C} - & \text{C} - & \text{C} - \text{H} \\ & & & \\ \text{H} & \text{H} & \text{H} & \text{H} \end{array}$ |
| 5. Pentane | C_5H_{12} | $\begin{array}{ccccc} \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \\ & & & & \\ \text{H} - \text{C} & - \text{C} - & \text{C} - & \text{C} - & \text{C} - \text{H} \\ & & & & \\ \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \end{array}$ |

Electron Dot Structure of Unsaturated Hydrocarbons

Ethene C_2H_4





Ethyne C_2H_2




| Name of Hydrocarbon | Molecular formula | Structural Formula |
|---------------------|-------------------|--|
| Alkenes : | | |
| 1. Ethene | C_2H_4 | $ \begin{array}{c} H & H \\ & \\ H - C = C - H \\ & \\ H & H \end{array} $ |
| 2. Propene | C_3H_6 | $ \begin{array}{c} H & H \\ & \\ H - C = C - C - H \\ & \\ H & H \end{array} $ |
| 3. Butene | C_4H_8 | $ \begin{array}{c} H & H \\ & \\ H - C = C - C - C - H \\ & & & \\ H & H & H & H \end{array} $ |
| Alkynes : | | |
| 1. Ethyne | C_2H_2 | $H - C \equiv C - H$ |
| 2. Propyne | C_3H_4 | $ \begin{array}{c} H \\ \\ H - C \equiv C - C - H \\ \\ H \end{array} $ |
| 3. Butyne | C_4H_6 | $ \begin{array}{c} H & H \\ & \\ H - C \equiv C - C - C - H \\ & \\ H & H \end{array} $ |

Carbon Compounds on the Basis of Structure


(i) Straight (unbranched) chain




(ii) Branched




Isomerism of C_5H_{12} (**Pentane**)



n-pentane




iso-pentane



Neo-pentane


These three above compounds has same molecular formula but different structures are called structural isomers and phenomenon is structural isomerism.

(iii) Cyclic





Cyclic Saturated

eg. C_6H_{12} (Hexane)



Cyclic unsaturated
 C_6H_6 (Benzene)



Functional Groups

- In hydrocarbon chain, one or more hydrogen atom is replaced by other atoms in accordance with their valencies. These are heteroatom.
- These heteroatom or group of atoms which make carbon compound reactive and decides its properties are called functional groups.

| Hetero atom | Functional group | Formula of functional group |
|-------------|---------------------|---|
| Cl/Br | Halo (Chloro/Bromo) | — Cl, — Br, — I |
| Oxygen | 1. Alcohol | — OH |
| | 2. Aldehyde | $\begin{array}{c} \text{H} \\ \diagup \\ \text{— C} \\ \diagdown \\ \text{O} \end{array}$ |
| | 3. Ketone | $\begin{array}{c} \text{— C —} \\ \\ \text{O} \end{array}$ |
| | 4. Carboxylic acid | $\begin{array}{c} \text{O} \\ \\ \text{— C — OH} \end{array}$ |
| Double bond | 1. Alkene group | $> \text{C} = \text{C} <$ |
| Triple bond | 2. Alkyne group | $\text{— C} \equiv \text{C —}$ |

Homologous Series

It is series of compounds in which the some functional group substitutes for the hydrogen in a carbon chain.

E.g., Alcohols – CH_3OH , $\text{C}_2\text{H}_5\text{OH}$, $\text{C}_3\text{H}_7\text{OH}$, $\text{C}_4\text{H}_9\text{OH}$

- Have same general formula.
- Any two homologues differ by $-\text{CH}_2$ group and difference in molecular mass is 14μ .
- Have same chemical properties but show gradual change in physical properties.

Nomenclature of Carbon Compounds

- Identify the number of carbon atoms in compounds.
- Functional group is indicated by suffix or prefix.

| Functional Group | Prefix/Suffix | Example |
|--------------------------|-----------------------------------|---|
| 1. Halogen | Prefix – Chloro, Bromo, Iodo etc. | $ \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H} - \text{C} - \text{C} - \text{C} - \text{Cl} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array} $ Chloro Propane |
| 2. Alcohol | Suffix – ol | $ \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H} - \text{C} - \text{C} - \text{C} - \text{OH} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array} $ Propanaol |
| 3. Aldehyde | Suffix – al | $ \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H} - \text{C} - \text{C} - \text{C} = \text{O} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array} $ Propanal |
| 4. Ketone | Suffix – one | $ \begin{array}{c} \text{H} \quad \quad \text{H} \\ \quad \quad \\ \text{H} - \text{C} - \text{C} - \text{C} - \text{H} \\ \quad \quad \\ \text{H} \quad \text{O} \quad \text{H} \end{array} $ Propanone |
| 5. Carboxylic acid | Suffix – oic acid | $ \begin{array}{c} \text{H} \quad \text{H} \quad \text{O} \\ \quad \quad \\ \text{H} - \text{C} - \text{C} - \text{C} - \text{OH} \\ \quad \\ \text{H} \quad \text{H} \end{array} $ Propanoic acid |
| 6. Alkene (– C = C –) | Suffix – ene | $ \begin{array}{c} \text{H} \quad \quad \text{H} \\ \quad \quad \\ \text{H} - \text{C} - \text{C} = \text{C} - \text{H} \\ \quad \quad \\ \text{H} \quad \quad \text{H} \end{array} $ Propene |

| | | |
|---------------------------------|--------------|--|
| 7. Alkyne ($-C \equiv C-$) | Suffix – yne | $ \begin{array}{c} \text{H} \\ \\ \text{H} - \text{C} - \text{C} \equiv \text{C} - \text{H} \\ \\ \text{H} \end{array} $ Propyne |
|---------------------------------|--------------|--|

Chemical Properties of Carbon Compounds


(a) Combustion




- Carbon and its compounds are used as fuels because they burn in air releasing lot of heat energy.
- Saturated hydrocarbon generally burn in air with blue and non-sooty flame.
- Unsaturated hydrocarbon burns in air with yellow sooty flame because percentage of carbon is higher than saturated hydrocarbon which does not get completely oxidized in air.

(b) Oxidation

Alcohols can be converted to carboxylic acid in presence of oxidizing agent alkaline KMnO_4 (potassium permanganate) or acidic potassium dichromate.




(c) Addition Reaction :



Unsaturated hydrocarbon add hydrogen in the presence of catalyst palladium or nickel. Vegetable oils are converted into vegetable ghee using this process. It is also called hydrogenation of vegetable oils.

(d) Substitution Reaction :



Important Carbon Compounds : Ethanol and Ethanoic acid


Physical Properties of Ethanol

- Colourless, pleasant smell and burning taste.

- Soluble in water.
- Volatile liquid with low boiling point of 351 K.
- Neutral compound.


Chemical Properties

(i) Reaction with Sodium :



This reaction is used as a test for ethanol by evolution of H_2 gas (Burn with pop sound).

(ii) Dehydration :




Physical Properties of Ethanoic acid


- Colourless liquid having sour taste and have smell of vinegar.
- Boiling point is 391 K.
- When pure CH_3COOH is freezed, it forms colourless ice like solid. So it is called glacial acetic acid.

Chemical Properties

(i) Esterification :




Sweet smelling ester is formed.




This is saponification as soap is prepared by this.

(ii) Reaction with base :



(iii) Reaction with carbonates and hydrogen carbonates :




Soaps and Detergents

- Soap is sodium or potassium salt of long chain carboxylic acid. *E.g.*, $\text{C}_{17}\text{H}_{35}\text{COO}^-\text{Na}^+$
- Soaps are effective only in soft water.
- Detergents are ammonium or sulphonate salt of long chain of carboxylic acid.
- Detergents are effective in both hard and soft water.

Soap molecule has :


- (i) Ionic (hydrophilic) part
- (ii) Long hydrocarbon chain (hydrophobic) part



Structure of soap molecule

Cleansing Action of Soap

- Most dirt is oily in nature and hydrophobic end attaches itself with dirt and the ionic end is surrounded with molecule of water. This results in formation of a radial structure called micelles.
- Soap micelles help to dissolve dirt and grease in water and cloth gets cleaned.



- The magnesium and calcium salt present in hard water react with soap molecule to form insoluble product called scum. This scum creates difficulty in cleansing action.
- By use of detergent, insoluble scum is not formed with hard water and cloths get cleaned effectively.

QUESTIONS

VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)

1. How does carbon attain noble gas configuration ?
2. Draw electron dot structure of water molecule.
3. Write the name and formula of 2nd member of homologous series having general formula C_nH_{2n} .
4. Name the first member of ketones.
5. What is glacial acetic acid ?
6. Why carbon is tetravalent ?
7. An organic compound burns with blue clear flame. Is it saturated or unsaturated compound ?
8. Write the molecular formula of ethanol.
9. Which of the following will show addition reaction : C_4H_{10} , C_2H_6 , C_2H_4 , CH_4 , C_3H_8 ?
10. Name the gas evolved when ethanoic acid is added to sodium carbonate ?
11. Write balanced chemical equation of dehydration of ethanol by hot conc. H_2SO_4 .
12. Name organic acid present in vinegar.
13. What is catenation ?
14. Why soap is not suitable for washing cloth when water is hard ?
15. How many covalent bonds are present in pentane (C_5H_{12}) ?

SHORT ANSWER TYPE QUESTIONS (2 Marks)

1. What are hydrocarbons ? Give examples.
2. Why does carbon atom forms a large number of compounds ?
3. Write down two characteristics of the compounds of an homologous series.
4. Covalent compounds generally don't conduct electricity. Why ?
5. Write down structural formula of :
 - (a) Propanone
 - (b) Hexanal
6. Why carbon is unique in nature ?

7. Which is better for health butter or vegetable oil ? Why ?
8. Complete the following reactions :
- (a) $\text{CH}_4 + \text{O}_2 \rightarrow$ (b) $\text{CH}_4 + \text{Cl}_2 \xrightarrow{\text{Sunlight}}$
9. Identify the functional group in following :
- (a) HCHO (b) CH_3COOH
(c) $\text{CH}_3\text{CH}_2\text{OH}$ (d) CH_3COCH_3
10. (a) Why is ethanol used in making of tincture iodine, cough syrup, tonic etc.
(b) What is the role of conc. H_2SO_4 in making ethane from ethanol ?

SHORT ANSWER TYPE QUESTIONS (3 Marks)

1. Differentiate between soap and detergents.
2. What is oxidizing agent ? Give two examples.
3. What is hydrogenation ? Write its industrial application.
4. What is homologous series ? Explain with the help of example.
5. Write IUPAC names of :
(a) $\text{HC} \equiv \text{CH}$ (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
(c) CH_2CHO
6. What is structural isomerism ? Draw isomers of pentane (C_5H_{12}).
7. A boy sharpens a pencil at both the ends and connects them to the poles of the battery. Will the current flow through the circuit ? Give reason.
8. A neutral organic compound is warmed with some ethanoic acid and a little of conc. H_2SO_4 to form ester, vapours having sweet smell are evolved. Write the chemical equation and what type of functional group is present in this organic compound ?

LONG ANSWER TYPE QUESTIONS (5 Marks)

1. Explain the cleansing action of soap with the help of diagram.
2. When ethanoic acid reacts with sodium hydrogen carbonate, the salt 'X' is formed and gas 'Y' is evolved.
(a) Identify 'X' and 'Y'.
(b) Write balanced chemical equation of above reaction.
(c) Describe a test to identify the gas 'Y' evolved.

Hints to Long Answer Type Questions

1. Page No. 74, Fig. 4.12 (Diagram of formation of micelles) of NCERT.
2. $2\text{CH}_3\text{COOH} + \text{Na}_2\text{CO}_3 \rightarrow 2\text{CH}_3\text{COONa} + \text{H}_2\text{O} + \text{CO}_2$

